

Preliminary Exam - February, 2018

Real Analysis

- 1) a) Let $f : X \rightarrow \mathbb{R}$ be integrable, where (X, μ) is a measure space. Show that the set $S = \{x : f(x) \neq 0\}$ is of σ -finite measure.

b) Evaluate $\lim_{k \rightarrow \infty} \sum_{n=1}^{\infty} e^{-n^2/k}$

Justify your answer using measure theory theorems.

- 2) a) Let $f \in L_p[0, 1]$ and let $\epsilon > 0$. Show that if $p \geq 1$ then

$$m(\{x \in [0, 1] : |f(x)| \geq \epsilon\}) \leq \epsilon^{-p} \int |f|^p dm$$

- b) Let $f_n \in L_p[0, 1] \forall n$. Show that if

$$\lim_{n \rightarrow \infty} \|f_n - f\|_p = 0 \quad \text{holds in } L_p[0, 1] \quad \text{then } (f_n) \text{ converges in measure to } f.$$

- 3) a) Show that if $f \in L_2 \cap L_4$ then $f \in L_3$ holds.

- b) Let $1 \leq p < q < \infty$ Show $L_q[0, 1] \subset L_p[0, 1]$ holds.

- 4) a) State the Fubini Theorem

b) Using $f(x, y) = \begin{cases} 2 - 2^{-x} & \text{if } x = y \\ -2 + 2^{-x} & \text{if } x = y + 1 \\ 0 & \text{otherwise} \end{cases}$

where $f : X \times Y \rightarrow \mathbb{R}$, $X = Y = \mathbb{N}$ both equipped with counting measure, show that integrability of f in Fubini's Theorem cannot be removed from the hypothesis.