

METU - Department of Mathematics  
Graduate Preliminary Exam

Algebra I

February, 2009

**Duration:** 180 min.

1. Given a group  $G$ , we can construct a chain  $G_1 \xrightarrow{\pi_1} G_2 \xrightarrow{\pi_2} G_3 \xrightarrow{\pi_3} G_4 \xrightarrow{\pi_4} \dots$ , where  $G_1 = G$  and  $G_{n+1} = \text{Aut}(G_n)$ , and  $\pi_n(g)(x) = gxg^{-1}$  for all  $g$  and  $x$  in  $G_n$ , for all positive integers  $n$ .

- a) Show  $\pi_n(G_n) \trianglelefteq G_{n+1}$  for all positive integers  $n$ .
  - b) Assuming  $C(G) = \langle 1 \rangle$ , show that  $\pi_n$  is injective, and  $C_{G_{n+1}}(\pi_n(G_n)) = \langle 1 \rangle$  for all positive integers  $n$ .
- 

2. Let  $G$  be a finite group and  $p$  be the smallest prime divisor of  $|G|$ . Prove that if  $H$  is a subgroup of index  $p$  in  $G$ , then  $H \trianglelefteq G$ .

---

3. Let  $\Omega$  be a set, and for each  $i$  in  $\Omega$ , let  $K_i$  be a field. Then let  $R$  be the ring  $\prod_{i \in \Omega} K_i$ . Let  $\mathfrak{m}$  be a maximal ideal of  $R$ . If  $x$  is an element  $(x_i : i \in \Omega)$  of  $R$ , define  $S(x) = \{i \in \Omega : x_i \neq 0\}$ .

- a) Show that, for all  $x$  and  $y$  in  $R$ , if  $\Omega$  is the *disjoint* union of  $S(x)$  and  $S(y)$ , then exactly one of  $x$  and  $y$  is in  $\mathfrak{m}$ .
  - b) Show that the homomorphism  $x \mapsto x/1$  from  $R$  to the localization  $R_{\mathfrak{m}}$  is surjective.
  - c) Find the kernel of the homomorphism in (b).
  - d) What kind of ring is  $R_{\mathfrak{m}}$ ?
- 

4. Let  $R$  be a ring and  $e$  be an idempotent in  $R$ , that is,  $e^2 = e \neq 0$ .
- a) Show that  $eRe$  is a subring of  $R$  and  $e$  is the identity of  $eRe$ .
  - b) Show that if  $R$  is finite and contains no nonzero nilpotent elements, then we have  $eRe = eR$ .